




PRELIMINARY GUIDE TO THE IDENTIFICATION OF THE EARLY LIFE  
HISTORY STAGES OF CIRRHITID FISHES OF THE WESTERN CENTRAL  
NORTH ATLANTIC

BY

MARIA M. CRIALES



U.S. DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Southeast Fisheries Science Center  
Miami, FL 33149

April 2003



PRELIMINARY GUIDE TO THE IDENTIFICATION OF THE EARLY LIFE  
HISTORY STAGES OF CIRRHITID FISHES OF THE WESTERN CENTRAL  
NORTH ATLANTIC

BY

M. M. CRIALES

University of Miami  
Rosensteil School of Marine and Atmospheric Sciences  
4600 Rickenbacker Causeway  
Miami, FL 33149

U.S. DEPARTMENT OF COMMERCE  
Donald L. Evans, Secretary

National Oceanic and Atmospheric Administration  
Conrad C. Lautenbacher, Jr., Under Secretary for Oceans and Atmosphere

National Marine Fisheries Service  
William T. Hogarth, Assistant Administrator for Fisheries

April 2003

This Technical Memorandum series is used for documentation and timely communication of preliminary results, interim reports, or similar special-purpose information. Although the memoranda are not subject to complete formal review, editorial control, or detailed editing, they are expected to reflect sound professional work.

## NOTICE

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or material mentioned in this publication. No reference shall be made to NMFS or to this publication furnished by NMFS, in any advertising or sales promotion which would imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein or which has as its purpose any intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

This report should be cited as follows:

Criales, M. M. 2003. Preliminary guide to the identification of the early life history stages of cirrhitid fishes of the western central North Atlantic. NOAA Technical Memorandum NMFS-SEFSC-497, 4 p.

W. J. Richards, Editor. NOAA Fisheries, 75 Virginia Beach Drive, Miami, FL

This report will be posted on the Bethune Cookman College NOAA Cooperative web site later in 2003 at <[www4.cookman.edu/noaa/](http://www4.cookman.edu/noaa/)> and also appear on the SEFSC web site at <[www.sefsc.noaa.gov/](http://www.sefsc.noaa.gov/)>

It will be a chapter entitled Cirrhitidae in “The early life history stages of fishes of the western central North Atlantic”.

Copies may be obtained by writing:

the author at  
University of Miami  
Rosensteil School of Marine and Atmospheric Sciences  
4600 Rickenbacker Causeway  
Miami, FL 33149  
or

National Technical Information Center  
5825 Port Royal Road  
Springfield, VA 22161  
(800) 553-6847 or (703) 605-6000  
<<http://www.ntis.gov/numbers.htm>>

Species of the family Cirrhitidae are known in the western and eastern Atlantic, Indian, and Pacific, with the majority represented in the Indo-Pacific. All 35 species are found in tropical, coastal waters and are usually small and brightly colored (Robins et al. 1986). Cirrhitid fishes live in rocky and coral habitats (mostly in crevices) and have many superficial features in common with the scorpaenids (Böhlke & Chaplin 1968). Only one species, *Amblycirrhitus pinos*, is known from the central western North Atlantic.

*Amblycirrhitus pinos*, the redspotted hawkfish, has 41 to 44 lateral-line scales, with a tuft of cirri from a membrane near the tip of each dorsal spine. The body has five broad dark bars, the first three yellowish brown, the upper rounded part of the fourth black, and the fifth (across the caudal peduncle) black with white interspaces between the first four dark bars bisected by narrow yellowish brown bars (Randall 1968). The anterior part of the head and body and dorsal fin has bright orange-red spots. Adult specimens reach up to 8 cm (4 inches). The redspotted hawkfish has been reported in south Florida, Bahamas, Texas, northern South America, and St. Helena Island (Robins et al. 1986).

Information about cirrhitid larvae for the western Atlantic is very scarce. A drawing of a postflexion *A. pinos* larvae was provided by Johnson (1984) (Fig. 1B), but no descriptive notes accompanied it. Larvae of at least three Indo-Pacific species have been reared in laboratory (Tanaka & Suzuki 1991, Tanaka 1994, 1995) and other larvae have been described from plankton samples (e.g. Leis and Rennis 1983, Watson 1996a). Based on these descriptions and material examined from the CALCOFI study area, Watson (1996a) provided an excellent review of

cirrhitid larvae for the Indo-Pacific with general features to family level.

Cirrhitid larvae were rare in our plankton collections. Four examined specimens were captured in the Straits of Florida (all postflexion, 8.0-13.2 mm) and two others were obtained in proximity to Barbados (one preflexion, 4.2 mm and one postflexion, 12.3 mm). These specimens allow us to make a general description of hawkfish larvae. Illustrations (see Fig. 1) are presented for preflexion and postflexion stages and Table 1 provides meristic data.

Atlantic hawkfish larvae have a moderately elongate and slender body; the snout is acute, and the anterior border of the lower jaw ends in a pigmented barbel, which is larger in the postflexion stage and considered as the diagnostic family feature. The eyes are large with a relatively wide oval shape and well pigmented in the postflexion stage but eyes in the preflexion specimen were missing. The snout in the preflexion specimen is elongate and pointed and becomes shorter and less elongate later in development. The preflexion specimen shows the caudal, dorsal and anal fin anlage, with the caudal fin anlage surrounded by numerous thick striations. The gut in the preflexion is long about half of the body length. The postflexion stage has a bifurcate caudal fin. There are 26 myomeres in both preflexion and postflexion stages. Two pigmentation patterns were common on cirrhitid specimens. Both (preflexion and postflexion) possess four rounded brownish melanophores located dorsally on the midbrain and distributed in two paired rows, with the two posterior melanophores being larger. The other common pigmentation pattern is a row of paired melanophores (postflexion 8, preflexion 10) distributed along the lower mandible and continuing to


the anterior barbel. Besides these two common patterns, the preflexion specimen shows a group of melanophores running ventrally from the base of the pectoral fin to the end of the caudal fin anlage, and another group running dorsally along the dorsal fin anlage. The gut of the preflexion specimen was also well pigmented all along the ventral margin of the tail.

**Table Cirrhitidae 1. Meristics  
for the family Cirrhitidae**


Stage	Collection Site	Size(mm)	D <sub>1</sub>	D <sub>2</sub>	A	P <sub>1</sub>	P <sub>2</sub>	C	Myomere
<b>Preflexion</b>									
	Barbados	4.2							10+16
<b>Postflexion</b>									
	Starits of Florida	8.0	VIII?	12	9				
	Starits of Florida	8.6	IX	12	9				
	Starits of Florida	10.8	IX	12	9				
	Barbados	12.3	IX	12	8	15	I,5	12/13	26+
	Starits of Florida	13.2	X?	13?	damaged	15?	I,5	8/7	
<b>Adult</b>			X	11	III,6	14	I,5	8/7	10+16=26

Figure 1 Larvae of the cirrhitid fish *Amblycirrhitus pinos*. A). preflexion stage, specimen from Barbados. 4.2 mm NL. B) postflexion stage, 13.2 mm SL (from Johnson 1984).

A



B



### Literature Cited

- Bohlke, J. E. & C. C. G. Chaplin. 1968. Fishes of the Bahamas and adjacent tropical waters. Livingston, Wynnewood. 771 p.
- Johnson, D. G. 1984. Percoidei: development and relationships. Pages 464-498 *in* Ontogeny and systematics of fishes. H. G. Moser et al. (eds.). Amer. Soc. Ichthyol. Herpetol. Spec. Publ. (1): 760 p.
- Leis, J. M. & D. S. Rennis. 1983. The larvae of Indo-Pacific coral reef fishes. University of Hawaii Press. 269 p.
- Randall, J. E. 1968. Caribbean reef fishes. T. F. H. Publ., Neptune City, N. J. 318 p.
- Robins, C. R. & G. C. Ray. 1986. A field guide to Atlantic coast fishes of North America. The Peterson Field Guide series: (32). Houghton Mifflin Co., Boston, Mass. 354 p.
- Tanaka, Y. and K. Suzuki. 1991. Spawning, eggs and larvae of the hawfish, *Cirrhitichthys aurens* in an aquarium. Jpn. J. Ichthyol. 38: 283-288.
- \_\_\_\_\_. 1994. Reproductive behavior, eggs and larval development of two species of the *Cirrhitichthys*, *Cirrhitidae*- *c. oxycephalus* and *C. falco*. J. Fac. Mar. Sci. Tech. Tokai Univ. 38: 213-232 (In Japanese).
- \_\_\_\_\_. 1995. Reproductive behavior, eggs and larval development of three species, three genus of the *Cirrhitidae*. J. Fac. Mar. Sci. Tech. Tokai Univ. 39: 173-194 (In Japanese).
- Watson, W. 1996a. *Cirrhitidae*. Pages 1064-1067 *in* The early stages of fishes in the California Current Region. H. G. Moser (ed.). CalCOFI Atlas 33: 1505 p.